



Effects of Leaf Maturity and Solvent Extract on the Antioxidant Activity of *Litsea elliptica*

Harlinda KUSPRADINI^{1,2,3,†} · Maulidia Shufwatul MALA¹ · Agmi Sinta PUTRI^{1,2,3} · Najmia Afifah ZULFA¹ · Hayatus SA'ADAH⁴ · KISWANTO¹

ABSTRACT

Litsea elliptica, a Southeast Asian tree with a rich history of medicinal applications, is attracting increasing research attention. This study investigated the effects of leaf maturity and solvent selection on the extraction of bioactive compounds from *L. elliptica* leaves, specifically with regard to their antioxidant activity. 2,2'-Azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS) method was employed to quantify the free radical scavenging capacity of *L. elliptica* leaf extracts prepared using three different solvents (n-hexane, ethyl acetate, and ethanol) at three different leaf stages (tender, immature, and mature). These results highlight the significant effects of leaf maturity and solvent selection on the extraction of phenolic compounds and flavonoids from *L. elliptica* leaves. Ethanol is the most effective solvent for the extraction of bioactive compounds, particularly from mature leaves. The ethanol extraction of tender leaves demonstrated potential for optimizing the antioxidant content. Further research is necessary to investigate the specific factors influencing the observed differences in antioxidant activity between leaves of varying ages and the potential impacts of other bioactive compounds present in the leaves.

Keywords: *Litsea elliptica*, leaves maturity, extracts, solvent, antioxidant, 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS) method

1. INTRODUCTION

The genus *Litsea*, with its diverse range of therapeutically promising plants, has attracted considerable attention because of its wide range of functionalities. Several pharmacological studies have confirmed the traditional medicinal uses of *Litsea* species. A thorough and systematic investigation of the phytochemical con-

stituents and pharmacological properties of *Litsea*, especially its mechanisms of action and toxicology, is crucial for understanding its ethnomedicinal uses, exploring its therapeutic potential, and supporting the development of future healthcare products (Wang *et al.*, 2016). Among the most promising *Litsea* species, *Litsea elliptica* Blume, a prominent tropical tree native to Southeast Asia, has a long history of medicinal applications. Kuspradini *et*

Date Received March 23, 2024; Date Revised April 8, 2024; Date Accepted June 10, 2024; Published September 25, 2024

¹ Faculty of Forestry, Universitas Mulawarman, Samarinda 75123, Indonesia

² Research Center of Medicine and Cosmetics from Tropical Rainforest, Universitas Mulawarman, Samarinda 75123, Indonesia

³ Research Collaboration Center for Biomass-Based Nano Cosmetic, Collaboration of Mulawarman University and BRIN, Samarinda 75123, Indonesia

⁴ Department of Pharmaceutical Technology, Sekolah Tinggi Ilmu Kesehatan Samarinda, East Kalimantan 75242, Indonesia

[†] Corresponding author: Harlinda KUSPRADINI (e-mail: hkuspradini@fahutan.unmul.ac.id, <https://orcid.org/0000-0001-5716-5955>)

© Copyright 2024 The Korean Society of Wood Science & Technology. This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

al. (2018) reported that *L. elliptica* and *L. angulata* are the most dominant *Litsea* species in East Kalimantan, Indonesia. Seven diverse compounds, including flavonoids, alkaloids, and steroids, were previously found in the leaves and wood of *L. elliptica* extracts (Phoopha *et al.*, 2020). Three major compounds, 2-undecanol (36.35%), terpineol (30.52%), and 9-decen-2-ol (22.43%), were obtained from the essential oil of *L. elliptica* (Kuspradini *et al.*, 2021). This rich chemical composition contributes to the potential of plants for various applications in medicine and natural product research.

The leaves are particularly revered because of their therapeutic properties against a variety of ailments. Traditional practices often involve the application of crushed leaves to the forehead for headache relief, thereby demonstrating the diverse contributions of the plant to local healthcare. Beyond headaches, *L. elliptica* leaves are also used to treat stomach ulcers, fever, and asthma, and even repel insects, solidifying their valuable role in healthcare traditions in the region (Goh *et al.*, 2022; Grosvenor *et al.*, 1995; Jiwajinda *et al.*, 2002; Yang *et al.*, 2022a). The young leaves are consumed as a vegetable side dish and used as a flavoring material in Thai 'Nam Prik,' a local spicy dip (Ngernsaengsaruy *et al.*, 2011).

In addition to its ecological importance, *L. elliptica* has garnered attention for its diverse array of bioactive compounds and potential health benefits, including its antioxidant properties (Wong *et al.*, 2014). The discovery of new and safe antioxidants from natural sources holds immense potential for various applications, including natural antioxidants, functional foods, and nutraceuticals. This growing interest has fueled the exploration of plant-based sources of antioxidants, and phytochemical screening has emerged as a valuable method for identifying and investigating these bioactive compounds in plants (Lee *et al.*, 2020; Nkogo *et al.*, 2022). Understanding the chemical composition and antioxidant activities at various ripening stages can assist producers in deter-

mining the optimal time for harvesting plants (Um *et al.*, 2020; Yang *et al.*, 2022b). Optimizing the conditions and design parameters is crucial for scaling up the extraction processes. This involves considering factors such as solvent type, particle size, solid-solvent ratio, and temperature, as previously reported by Mindaryani *et al.* (2023) and Mun *et al.* (2020).

Previous studies have highlighted the presence of various phenolic compounds in *L. elliptica*, including flavonoids and tannins, which possess antioxidant properties. However, the influence of leaf maturity and solvent extracts on antioxidant activity remains largely unexplored. This study aimed to address this gap by investigating the variation in the antioxidant activity of *L. elliptica* leaves and exploring the influence of leaf stage (tender, immature, and mature) and solvent extracts (n-hexane, ethyl acetate, and ethanol) on the antioxidant activity of the extracts. The 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS) method, a widely recognized and sensitive antioxidant assay, was used in this study to quantify the free radical scavenging capacity of different extract types.

Harvesting *L. elliptica* leaves at the best stage and extracting them using an appropriate solvent is crucial for enhancing the potential of this plant as a natural source of antioxidants for therapeutic and health-promoting applications. Comprehensive analysis of these factors can provide valuable insights into effective and sustainable extraction strategies and maximize the potential of *L. elliptica* leaves as a natural source of antioxidants for therapeutic and health-promoting applications.

2. MATERIALS and METHODS

2.1. Sample collection

This study was conducted at the Laboratory of Forest Product Chemistry and Renewable Energy and the Laboratory of Forest Cultivation, Faculty of Forestry,

Mulawarman University, Samarinda, Brazil. *Litsea elliptica* leaves were obtained from 6-month-old plants grown in the same faculty. Leaf samples were collected at three stages of maturity: tender, immature, and mature.

2.2. Extraction

After air-drying, the leaf samples were ground into a powder using a mechanical grinder. The extract preparation method was adopted from Kuspradini *et al.* (2023). Subsequently, maceration was performed using n-hexane, ethyl acetate, and 96% ethanol (technical grade) to extract the powder (10 g). The powder was first extracted with n-hexane for 24 h at a 1:1 (w/v) ratio, and the resulting solution was filtered using filter paper (with a thickness and basic weight specification of 0.0893 m³ and 80 ± 5 g, respectively). Additional extraction was performed with ethyl acetate by soaking the residue in the solvent at a 1:1 (w/v) ratio for 24 h to produce an ethyl acetate extract from the filtrate. Finally, the residue was extracted with ethanol at a 1:1 (w/v) ratio for 24 h, and the resulting filtrate was collected as the ethanol extract. All extractions were performed on a shaker. The filtrate was evaporated using a dehydration machine at 40°C to obtain a crude extract. All extracts (n-hexane, ethyl acetate, and ethanol extracts) were then analyzed using antioxidant assays for total phenolic and flavonoid content.

2.3. Total phenolic content

The phenolic content of the *L. elliptica* leaf extract was determined according to a previously reported method, with a few modifications (Kuspradini *et al.*, 2016). Briefly, the sample of 100 µL (1 mg of extract in 10 mL aquadest) was mixed with 400 µL aquadest, 250 µL Folin-Ciocalteu reagent, and 1,250 µL of 7.5% sodium carbonate. After 1 h of incubation, the absorbance was measured at 760 nm using a spectrophotometer. Total

phenolic content (TPC) was measured using a gallic acid calibration curve and expressed in milligrams of gallic acid equivalent (mg GAE).

2.4. Total flavonoid content

A spectrophotometric method (Kuspradini *et al.*, 2016) using a catechin standard was employed with minor modifications. A mixture was prepared by combining 100 µL extract (1 mg/10 mL) with 700 µL aquadest, 100 µL sodium nitrite, 100 µL aluminum chloride, and 500 µL sodium hydroxide. The homogenized mixture was left undisturbed in the dark for 10 min at room temperature. Absorbance was measured at 510 nm using a spectrophotometer. The flavonoid content was calculated using the calibration curve of the catechin standard and presented as milligram catechin equivalents (mg CE).

2.5. 2,2'-Azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid assay

Antioxidant activity against ABTS radicals was measured according to a previously reported protocol (Toppo *et al.*, 2019). The ABTS solution was prepared by combining a 7.4 mM ABTS solution with a 2.6 mM potassium persulphate solution in a 2:1 ratio. The mixture was incubated for 16 h at room temperature in the dark. The solution was then combined with 1 mL of ABTS solution and 60 mL of methanol. The absorbance of (0.70 ± 0.02) units at a wavelength of 734 nm was measured using a spectrophotometer. Different concentrations (6.25, 12.5, 25, 50, and 100 ppm) of 100 µL of extract were mixed with 900 µL ABTS solution and shaken for 45 s. After 15 min of incubation in the dark, the absorbance of the resulting mixture was measured at 734 nm using a spectrophotometer. Antioxidants were quantified as the percentage of ABTS free-radical scavenging activity. Ascorbic acid (vitamin C) was used

as a positive control. The scavenging activity was calculated using the following formula:

$$\text{Percentage (\%)} \text{ of scavenging activity} = \frac{[(\text{Absorbance of control} - \text{Absorbance of sample}) / \text{Absorbance of control}] \times 100}{(1)}$$

2.6. Data analysis

The TPC and total flavonoid content (TFC) were calculated using equations based on the linear regression models of gallic acid and catechin, respectively. The models used quercetin concentration as the independent variable (abscissa) and the average absorbance of triplicate measurements at each concentration as the dependent variable (ordinate).

The antioxidant capacity is expressed as the IC50 value, which represents the concentration of the extract required to inhibit 50% of ABTS free radicals. IC50 was calculated using a third-order polynomial regression equation, with the series of extract concentrations and the percentage of ABTS free-radical scavenging activity as the independent (abscissa) and dependent (ordinate) variables, respectively.

3. RESULTS and DISCUSSION

In this study the TPC of *L. elliptica* leaves showed a significant increase with leaf maturity for all solvent extracts (Fig. 1), similar to report by Wong *et al.* (2014) whereas, Nobossé *et al.* (2018) reported a similar increase in *Moringa oleifera* L. leaves, except for the aqueous extract. These findings suggest that the accumulation of phenolic compounds in leaves is a common phenomenon during plant growth and maturation.

According to Fig. 1 the TPC content of n-hexane extracts, ranged from 0.159 ± 0.003 mg to 0.226 ± 0.003 mg GAE/g extract for tender leaves and 0.194 ± 0.012 mg GAE/g extract for immature leaves to 0.357 ± 0.003 mg GAE/g extract for mature leaves. Similar trends were observed in ethyl acetate and ethanol extracts, with contents ranging from 0.343 ± 0.011 to 0.357 ± 0.03 mg GAE/g extract, 0.343 ± 0.011 to 0.357 ± 0.003 mg GAE/g extract, and 0.307 ± 0.005 to 0.568 ± 0.006 mg GAE/g extract, for tender, immature, and mature leaves, respectively. The results indicated that the TPC in ethanol was substantially higher than that in ethyl acetate and n-hexane. The choice of solvent can significantly affect the extraction yield of phenolic compounds.




Fig. 1. Total phenolic content (TPC) and total flavonoid content (TFC) of *Litsea elliptica* extracts. GAE: gallic acid equivalent, CE: catechin equivalents.

This highlights the importance of optimizing the solvent selection to maximize the extraction of the desired bioactive compounds. Variations in the extractable total phenol and flavonoid contents between different *Litsea* species, including *L. elliptica*, have been reported (Salehi *et al.*, 2019). This suggests that the specific composition and concentration of phenolic compounds may vary depending on the *Litsea* species and other factors, such as environmental conditions. The results indicated that the TPC in ethanol was substantially higher than that in ethyl acetate and n-hexane. A separate study (Goh *et al.*, 2022) found that young *L. elliptica* leaves yielded the most phenolic compounds when extracted with methanol compared to other solvents. This observation confirmed the effectiveness of polar solvents for extracting phenolic compounds from *L. elliptica* leaves.

The data presented in Fig. 1 show that the TFC in *L. elliptica* leaves exhibited complex trends depending on both the leaf maturity stage and the solvent extract. This finding highlights the significant interplay between these two factors during flavonoid extraction. Although the TFC increased slightly in the n-hexane extracts with increasing maturity, the difference was minimal. Conversely, the TFC of the ethyl acetate and ethanol extracts decreased from tender to mature leaves, similar to the findings in *Morus alba*. These findings suggest that the

relationship between leaf maturity and flavonoid content may not be universal and may vary depending on the plant species and extraction conditions.

Remarkably, unlike TPC, TFC levels were not dependent on solvent polarity, highlighting the distinct factors that influence the extraction of these two classes of bioactive compounds. Overall, these data reveal the importance of considering both leaf maturity and solvent choice when optimizing extraction protocols to maximize the yield of flavonoids from *L. elliptica* leaves.

The third-order polynomial regression model accurately predicted the relationship between the concentration and antioxidant activity of *L. elliptica*, as shown in Fig. 2. The strong correlation coefficient ($R^2 = 0.9825$ to 1) indicated a direct and significant relationship between the extract concentration and antioxidant activity measured by the ABTS assay. This suggests that increasing the concentration of the extract resulted in a proportional increase in antioxidant activity.

IC50, a widely employed parameter for measuring antioxidant activity, signifies the concentration at which a substance, such as ABTS, inhibits 50% of the free radical activity. Lower IC50 values indicate stronger antioxidant activity. The overall findings in this study suggest that the ethanolic extract from *L. elliptica* leaves has antioxidant properties ranging from active to highly




Fig. 2. Relationship between *Litsea elliptica* extract concentration and antioxidant activity measured by ABTS assay. ABTS: 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid.

active (15.34–6.35 ppm). According to Moga’s criteria, IC50 values $\leq 10 \mu\text{g/mL}$ are highly active, 10–150 $\mu\text{g/mL}$ are active, 150–500 $\mu\text{g/mL}$ are moderately active, and $> 500 \mu\text{g/mL}$ are inactive (Table 1; Moga *et al.*, 2021).

Highlighting the importance of solvent polarity, the data reveals that ethanol, a highly polar solvent, exhibits the strongest antioxidant activity (Purwanto *et al.*, 2017). This reinforces the observed trend of increasing activity with increasing solvent polarity (Herrera-Pool *et al.*, 2021; Huh *et al.*, 2022).

This emphasizes the importance of selecting an appropriate solvent to maximize the extraction of the desired bioactive compounds, similar to a study that showed appreciably high antioxidant potential of methanolic extracts of *L. elliptica* leaves (Suksamerkun *et al.*, 2013).

The effect of leaf maturity on antioxidant activity varied across the different solvent extracts. Although tender leaves extracted with ethanol displayed the highest activity, n-hexane and ethyl acetate extracts showed relatively consistent activity across all maturity stages. This suggests that to maximize the antioxidant content, harvesting *L. elliptica* leaves at the tender stage might be best when using ethanol as the extraction solvent. These findings highlight the potential of *L. elliptica*

leaves as a rich source of natural antioxidants, particularly when extracted with ethanol. The strong correlation between concentration and activity, coupled with the high accuracy of the regression model, provides valuable tools for optimizing the extraction process.

Additionally, a significant correlation was observed between ABTS radical-scavenging activity and leaf age in *L. elliptica*, with younger leaves showing higher ABTS radical-scavenging activity than older leaves. A similar link was observed in a previous study on *A. arguta* accessions (Tan *et al.*, 2021).

However, it is important to acknowledge the complex interplay between the factors that influence antioxidant activity. Although a positive correlation between antioxidant activity and total phenol/flavonoid content has been reported (Faiku *et al.*, 2019; Koraqi and Lluga-Rizani, 2022), this relationship is not always straightforward. Factors such as synergistic and antagonistic interactions between different types of phenols and flavonoids, the presence of other bioactive compounds, and the specificity of the ABTS method can contribute to this complexity (Giada, 2013; Hidalgo *et al.*, 2010; Kadum *et al.*, 2019; Munteanu and Apetrei, 2021). Therefore, a comprehensive understanding of these factors is crucial for accurately interpreting the relationship between ABTS activity and phenolic/flavonoid content. A high phenolic/

Table 1. Antioxidant activity criteria of *Litsea elliptica* leaves based on the IC50

Extract	Activity	Tender	Immature	Mature
n-Hexane	IC50 (ppm)	123.36	133.31	137.24
	Criteria	Moderately active	Moderately active	Moderately active
Ethyl acetate	IC50 (ppm)	11.13	15.42	10.78
	Criteria	Active	Active	Active
Ethanol	IC50 (ppm)	6.35	15.34	12.19
	Criteria	Highly active	Active	Active
Ascorbic acid	IC50 (ppm)	2.17		
	Criteria	Highly active		

flavonoid content alone cannot guarantee a strong antioxidant activity.

Additional investigations exploring the specific compounds responsible for the observed antioxidant activity and the potential applications of *L. elliptica* extracts in various fields are necessary to fully elucidate the potential of this promising natural resource.

4. CONCLUSIONS

The data presented in this study confirmed the influence of leaf maturity and solvent selection on the extraction of phenolic compounds and flavonoids from *L. elliptica* leaves. These findings underscore the importance of considering these factors when optimizing extraction protocols to maximize the yield of bioactive compounds with potential health benefits. These data suggest that ethanol is the most effective solvent for extracting phenols from mature *L. elliptica* leaves.

These results suggest that polar solvents are suitable for the extraction of antioxidant compounds from plant materials, particularly from *L. elliptica*. Furthermore, harvesting tender leaves may improve the antioxidant content when ethanol is used as the extraction solvent. It is important to note that the antioxidant activity of *L. elliptica* leaf extract can be influenced by antioxidant compounds other than phenols and flavonoids.

Further research is required to explore the specific factors contributing to the observed differences in antioxidant activity between leaves of different ages. This could involve analyzing the changes in individual phenol and flavonoid profiles and investigating the potential influence of other bioactive compounds present in the leaves.

CONFLICT of INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENT

The authors acknowledge the financial support provided by the Ministry of Education, Culture, Research, and Technology, Indonesia (grant number: 299/UN17.L1/HK/2023).

REFERENCES

- Faiku, F., Buqaj, L., Haziri, A. 2019. Phytochemicals and antioxidant study of *Teucrium chamaedrys* (L.) plant. *Agriculture and Forestry* 65(1): 137-145.
- Giada, M.L.R. 2013. Food Phenolic Compounds: Main Classes, Sources and their Antioxidant Power. In: *Oxidative Stress and Chronic Degenerative Diseases: A Role for Antioxidants*, Ed. by Morales-González, J.A. Instituto Politécnico Nacional, Mexico City, Mexico.
- Goh, M.P.Y., Kamaluddin, A.F., Tan, T.J.L., Yasin, H., Taha, H., Jama, A., Ahmad, N. 2022. An evaluation of the phytochemical composition, antioxidant and cytotoxicity of the leaves of *Litsea elliptica* Blume: An ethnomedicinal plant from Brunei Darussalam. *Saudi Journal of Biological Sciences* 29(1): 304-317.
- Grosvenor, P.W., Gothard, P.K., McWilliam, N.C., Supriono, A., Gray, D.O. 1995. Medicinal plants from Riau province, Sumatra, Indonesia. Part 1: Uses. *Journal of Ethnopharmacology* 45(2): 75-95.
- Herrera-Pool, E., Ramos-Díaz, A.L., Lizardi-Jiménez, M.A., Pech-Cohuo, S., Ayora-Talavera, T., Cuevas-Bernardino, J.C., García-Cruz, U., Pacheco, N. 2021. Effect of solvent polarity on the ultrasound assisted extraction and antioxidant activity of phenolic compounds from habanero pepper leaves (*Capsicum chinense*) and its identification by UPLC-PDA-ESI-MS/MS. *Ultrasonics Sonochemistry* 76: 105658.
- Hidalgo, M., Sánchez-Moreno, C., de Pascual-Teresa, S.

2010. Flavonoid–flavonoid interaction and its effect on their antioxidant activity. *Food Chemistry* 121(3): 691-696.
- Huh, J.S., Lee, S., Kim, D.S., Choi, M.S., Choi, H., Lee, K.H. 2022. Antioxidative and circadian rhythm regulation effect of *Quercus gilva* extract. *Journal of the Korean Wood Science and Technology* 50(5): 338-352.
- Jiwajinda, S., Santisopasri, V., Murakami, A., Kawanaka, M., Kawanaka, H., Gasquet, M., Eilas, R., Balansard, G., Ohigashi, H. 2002. *In vitro* anti-tumor promoting and anti-parasitic activities of the quassinoids from *Eurycoma longifolia*, a medicinal plant in Southeast Asia. *Journal of Ethnopharmacology* 82(1): 55-58.
- Kadum, H., Hamid, A.A., Abas, F., Ramli, N.S., Mohammed, A.K.S., Muhialdin, B.J., Jaafar, A.H. 2019. Bioactive compounds responsible for antioxidant activity of different varieties of date (*Phoenix dactylifera* L.) elucidated by 1H- NMR based metabolomics. *International Journal of Food Properties* 22(1): 462-476.
- Koraqi, H., Llugja-Rizani, K. 2022. Effect of extraction solvent on bioactive compounds and antioxidant activity of *Cichorium intybus* L. grown in Kosovo. *Agriculture and Forestry* 68(1): 159-171.
- Kuspradini, H., Putri, A.S., Diana, R. 2018. Potensi Tumbuhan Genus *Litsea*. Mulawarman University Press, Samarinda, Indonesia.
- Kuspradini, H., Rosiarto, A.M., Putri, A.S., Kusuma, I.W. 2016. Antioxidant and toxicity properties of anthocyanin extracted from red flower of four tropical shrubs. *Nusantara Bioscience* 8(2): 135-140.
- Kuspradini, H., Sinta, S., Silau, S., Putri, A.S. 2021. Karakteristik Minyak Atsiri Dari Tumbuhan Aromatik Hutan Tropis Jenis *Litsea* spp Dan Potensinya Sebagai Antimikroba. IPB Press, Bogor, Indonesia.
- Lee, M., Jeong, S.H., Mun, S.P. 2020. Conditions for the extraction of polyphenols from radiata pine (*Pinus radiata*) bark for bio-foam preparation. *Journal of the Korean Wood Science and Technology* 48(6): 861-868.
- Mindaryani, A., Sulton, A., Setiawan, F.A., Rahayuningsih, E. 2023. Natural dye extraction from Merbau (*Intsia bijuga*) sawdust: Optimization. *Journal of the Korean Wood Science and Technology* 51(6): 481-492.
- Moga, D.K., Adipo, N., Matu, E.N., Kirira, P.G. 2021. Antioxidant and antiproliferative activity of *Azadirachta indica* A. Juss extracts against cancer cell lines: An experimental study. *African Journal of Health Sciences* 34(5): 650-656.
- Mun, J.S., Kim, H.C., Mun, S.P. 2020. Chemical characterization of neutral extracts prepared by treating *Pinus radiata* bark with sodium bicarbonate. *Journal of the Korean Wood Science and Technology* 48(6): 878-887.
- Munteanu, I.G., Apetrei, C. 2021. Analytical methods used in determining antioxidant activity: A review. *International Journal of Molecular Sciences* 22(7): 3380.
- Nantitanon, W., Yotsawimonwat, S., Okonogi, S. 2010. Factors influencing antioxidant activities and total phenolic content of guava leaf extract. *LWT-Food Science and Technology* 43(7): 1095-1103.
- Ngernsaengsaruy, C., Middleton, D.J., Chayamarit, K. 2011. A revision of the genus *Litsea* Lam. (Lauraceae) in Thailand. *Thai Forest Bulletin (Botany)* 39: 40-119.
- Nkogo, L.F.E., Bopenga, C.S.A.B., Ngohang, F.E., Mengome, L.E., Angone, S.A., Engonga, P.E. 2022. Phytochemical and anti-termite efficiency study of *Guibourtia tessmanii* (harms) J. Léonard (Kévazingo) bark extracts from Gabon. *Journal of the Korean Wood Science and Technology* 50(2): 113-125.
- Nobossé, P., Fombang, E.N., Mbofung, C.M.F. 2018. Effects of age and extraction solvent on phytochemical content and antioxidant activity of fresh

- Moringa oleifera* L. leaves. Food Science & Nutrition 6(8): 2188-2198.
- Phoophu, S., Wattanapiromsakul, C., Pitakbut, T., Dej-adisai, S. 2020. Chemical constituents of *Litsea elliptica* and their alpha-glucosidase inhibition with molecular docking. Pharmacognosy Magazine 16(70): 327-334.
- Purwanto, D., Bahri, S., Ridhay, A. 2017. Uji aktivitas antioksidan ekstrak buah purnajiwa (*Kopsia arborea* Blume.) dengan berbagai pelarut. Kovalen 3(1): 24-32.
- Salehi, B., Ata, A., Kumar, N.V.A., Sharopov, F., Ramírez-Alarcón, K., Ruiz-Ortega, A. 2019. Anti-diabetic potential of medicinal plants and their active components. Biomolecules 9(10): 551.
- Suksamerkun, W., Thongsomchitt, S., Wongkrajang, Y., Tamsiririrkkul, R., Kitphati, W., Thongpraditchote, S. 2013. Screening of antioxidant activity of vegetables in Thailand. Journal of Asian Association of Schools of Pharmacy 2: 254-261.
- Tan, C., Wang, Z., Feng, X., Irfan, M., Changjiang, L. 2021. Identification of bioactive compounds in leaves and fruits of *Actinidia arguta* accessions from northeastern China and assessment of their antioxidant activity with a radical-scavenging effect. Biotechnology & Biotechnological Equipment 35(1): 593-607.
- Toppo, A.J., Chandra, S., Jha, D., Mazumder, P.M. 2019. *In vitro* evaluation of anti-acetylcholinesterase and free radical scavenging potential of leaf extracts of some selected medicinal plants. Asian Pacific Journal of Tropical Biomedicine 9(2): 60-65.
- Um, M., Kim, J.W., Lee, J.W. 2020. Optimization of ascorbic acid extraction from rugosa rose (*Rosa rugosa* Thunb.) fruit using response surface methodology and validation of the analytical method. Journal of the Korean Wood Science and Technology 48(3): 364-375.
- Wang, Y.S., Wen, Z.Q., Li, B.T., Zhang, H.B., Yang, J.H. 2016. Ethnobotany, phytochemistry, and pharmacology of the genus *Litsea*: An update. Journal of Ethnopharmacology 181: 66-107.
- Wong, M.H., Lim, L.F., bin Ahmad, F., bin Assim, Z. 2014. Antioxidant and antimicrobial properties of *Litsea elliptica* Blume and *Litsea resinosa* Blume (Lauraceae). Asian Pacific Journal of Tropical Biomedicine 4(5): 386-392.
- Yang, J., Choi, W.S., Lee, S.Y., Kim, M., Park, M.J. 2022a. Antioxidant activities of essential oils from *Citrus × natsudaidai* (Yu. Tanaka) Hayata peels at different ripening stage. Journal of the Korean Wood Science and Technology 50(4): 272-282.
- Yang, J., Lee, S.Y., Na, H., Jang, S.K., Park, M.J. 2022b. Evaluation of anti-asthmatic activity of essential oils from the Lauraceae family in lipopolysaccharide (LPS)-stimulated NCI-H292 cells. Journal of the Korean Wood Science and Technology 50(6): 414-426.